

Set Mathematics

MERT TOPALOĞLU 2 HAZ 2021, 19:37

SET MATHEMATICS

Set (mathematics)

From Wikipedia, the free encyclopedia

A set in mathematics is a collection of well defined and distinct objects, considered as an object in its own right. Sets are one of the most fundamental concepts in mathematics. Developed at the end of the 19th century, set theory is now a ubiquitous part of mathematics, and can be used as a foundation from which nearly all of mathematics can be derived. In mathematics education, elementary topics such as Venn diagrams are taught at a young age, while more advanced concepts are taught as part of a university degree.



The intersection of two sets is made up of the objects contained in both sets, shown in a Venn diagram.

Contents

- 1 Definition
- 2 Describing sets
- 3 Membership
 - 3.1 Subsets
 - 3.2 Power sets
- 4 Cardinality
- 5 Special sets
- 6 Basic operations
 - 6.1 Unions
 - 6.2 Intersections
 - 6.3 Complements
 - 6.4 Cartesian product
- 7 Applications
- 8 Axiomatic set theory
- 9 Principle of inclusion and exclusion
- 10 See also
- 11 Notes
- 12 References
- 13 External links

Definition

A set is a well defined collection of objects. Georg Cantor, the founder of set theory, gave the following definition of a set at the beginning of his *Beiträge zur Begründung der transfiniten Mengenlehre*:^[1]

A set is a gathering together into a whole of definite, distinct objects of our perception [Anschauung] and of our thought – which are called elements of the set.

The elements or members of a set can be anything: numbers, people, letters of the alphabet, other sets, and so on. Sets are conventionally denoted with capital letters. Sets *A* and *B* are equal if and only if they have precisely the same elements.^[2]

As discussed below, the definition given above turned out to be inadequate for formal mathematics; instead, the notion of a "set" is taken as an undefined primitive in axiomatic set theory, and its properties are defined by the Zermelo–Fraenkel axioms. The most basic properties are that a set "has" elements, and that two sets are equal (one and the same) if and only if every element of one is an element of the other.

Describing sets

There are two ways of describing, or specifying the members of, a set. One way is by intensional definition, using a rule or semantic description:

A is the set whose members are the first four positive integers.
B is the set of colors of the French flag.

The second way is by extension – that is, listing each member of the set. An extensional definition is denoted by enclosing the list of members in curly brackets:

C = {4, 2, 1, 3}
D = {blue, white, red}.

Every element of a set must be unique; no two members may be identical. (A multiset is a generalized concept of a set that relaxes this criterion.) All set operations preserve this property. The order in which the elements of a set or multiset are listed is irrelevant (unlike for a sequence or tuple). Combining these two ideas into an example

{6, 11} = {11, 6} = {11, 6, 6, 11}

because the extensional specification means merely that each of the elements listed is a member of the set.

For sets with many elements, the enumeration of members can be abbreviated. For instance, the set of the first thousand positive integers may be specified extensionally as:

{1, 2, 3, ..., 1000},

where the ellipsis ("...") indicates that the list continues in the obvious way. Ellipses may also be used where sets have infinitely many members. Thus the set of positive even numbers can be written as {2, 4, 6, 8, ... }.

The notation with braces may also be used in an intensional specification of a set. In this usage, the braces have the meaning "the set of all ...". So, *E* = {playing card suits} is the set whose four members are ♠, ♥, ♣, and ♦. A more general form of this is set-builder notation, through which, for instance, the set *F* of the twenty smallest integers that are four less than perfect squares can be denoted:

$$F = \{n^2 - 4 : n \text{ is an integer, and } 0 \leq n \leq 19\}.$$

In this notation, the colon (":") means "such that", and the description can be interpreted as "F is the set of all numbers of the form $n^2 - 4$, such that n is a whole number in the range from 0 to 19 inclusive." Sometimes the vertical bar (":") is used instead of the colon.

One often has the choice of specifying a set intensionally or extensionally. In the examples above, for instance, $A = C$ and $B = D$.

Membership

Main article: Element (mathematics)

The key relation between sets is *membership* – when one set is an element of another. If *a* is a member of *B*, this is denoted $a \in B$, while if *c* is not a member of *B* then $c \notin B$. For example, with respect to the sets $A = \{1,2,3,4\}$, $B = \{\text{blue, white, red}\}$, and $F = \{n^2 - 4 : n \text{ is an integer, and } 0 \leq n \leq 19\}$ defined above,

$4 \in A$ and $285 \in F$; but
 $9 \notin F$ and green $\notin B$.

Subsets

Main article: Subset

If every member of set *A* is also a member of set *B*, then *A* is said to be a *subset* of *B*, written $A \subseteq B$ (also pronounced *A is contained in B*). Equivalently, we can write $B \supseteq A$, read as *B is a superset of A*, *B includes A*, or *B contains A*. The relationship between sets established by \subseteq is called *inclusion* or *containment*.

If *A* is a subset of, but not equal to, *B*, then *A* is called a *proper subset* of *B*, written $A \subset B$ (*A is a proper subset of B*) or $B \supset A$ (*B is a proper superset of A*).

Note that the expressions $A \subset B$ and $B \supset A$ are used differently by different authors; some authors use them to mean the same as $A \subset B$ (respectively $B \supset A$), whereas other use them to mean the same as $A \subset B$ (respectively $B \supseteq A$).

Example:

- The set of all men is a proper subset of the set of all people.
- $\{1, 3\} \subset \{1, 2, 3, 4\}$.
- $\{1, 2, 3, 4\} \subset \{1, 2, 3, 4\}$.

The empty set is a subset of every set and every set is a subset of itself.

- $\emptyset \subseteq A$.
- $A \subseteq A$.

An obvious but useful identity, which can often be used to show that two seemingly different sets are equal:

- $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$.

A partition of a set *S* is a set of nonempty subsets of *S* such that every element *x* in *S* is in exactly one of these subsets.

Power sets


Main article: Power set

The power set of a set *S* is the set of all subsets of *S*, including *S* itself and the empty set. For example, the power set of the set {1, 2, 3} is { {}, {1, 2}, {1, 3}, {2, 3}, {1, 2, 3}, {1, 2, 3}, {} }. The power set of a set *S* usually written as *P*(*S*).

The power set of a finite set with *n* elements has 2^n elements. This relationship is one of the reasons for the terminology *power set*. For example, the set {1, 2, 3} contains three elements, and the power set shown above contains $2^3 = 8$ elements.

The power set of an infinite (either countable or uncountable) set is always uncountable. Moreover, the power set of a set is always strictly "bigger" than the original set in the sense that there is no way to pair the elements of a set *S* with the elements of its power set *P*(*S*) such that every element of *S* is paired with exactly one element of *P*(*S*), and every element of *P*(*S*) is paired with exactly one element of *S*. (There is never a bijection from *S* onto *P*(*S*)).

Every partition of a set *S* is a subset of the powerset of *S*.



Cardinality

Main article: Cardinality

The cardinality $|S|$ of a set S is "the number of members of S ." For example, if $B = \{\text{blue, white, red}\}$, $|B| = 3$.

There is a unique set with no members and zero cardinality, which is called the *empty set* (or the *null set*) and is denoted by the symbol \emptyset (other notations are used; see empty set). For example, the set of all three-sided squares has zero members and thus is the empty set. Though it may seem trivial, the empty set, like the number zero, is important in mathematics; indeed, the existence of this set is one of the fundamental concepts of axiomatic set theory.

Some sets have infinite cardinality. The set \mathbf{N} of natural numbers, for instance, is infinite. Some infinite cardinalities are greater than others. For instance, the set of real numbers has greater cardinality than the set of natural numbers. However, it can be shown that the cardinality of (which is to say, the number of points on) a straight line is the same as the cardinality of any segment of that line, of the entire plane, and indeed of any finite-dimensional Euclidean space.

Special sets

There are some sets which hold great mathematical importance and are referred to with such regularity that they have acquired special names and notational conventions to identify them. One of these is the empty set, denoted $\{\}$ or \emptyset . Another is the unit set $\{x\}$ which contains exactly one element, namely x .^[2] Many of these sets are represented using blackboard bold or bold typeface. Special sets of numbers include:

- \mathbf{P} or \mathbf{P} , denoting the set of all primes: $\mathbf{P} = \{2, 3, 5, 7, 11, 13, 17, \dots\}$.
- \mathbf{N} or \mathbf{N} , denoting the set of all natural numbers: $\mathbf{N} = \{1, 2, 3, \dots\}$ (sometimes defined containing 0).
- \mathbf{Z} or \mathbf{Z} , denoting the set of all integers (whether positive, negative or zero): $\mathbf{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$.
- \mathbf{Q} or \mathbf{Q} , denoting the set of all rational numbers (that is, the set of all proper and improper fractions): $\mathbf{Q} = \{a/b : a, b \in \mathbf{Z}, b \neq 0\}$. For example, $1/4 \in \mathbf{Q}$ and $11/6 \in \mathbf{Q}$. All integers are in this set since every integer a can be expressed as the fraction $a/1$ ($\mathbf{Z} \subseteq \mathbf{Q}$).
- \mathbf{R} or \mathbf{R} , denoting the set of all real numbers. This set includes all rational numbers, together with all irrational numbers (that is, numbers which cannot be rewritten as fractions, such as $\sqrt{2}$, as well as transcendental numbers such as π , e and numbers that cannot be defined).
- \mathbf{C} or \mathbf{C} , denoting the set of all complex numbers: $\mathbf{C} = \{a + bi : a, b \in \mathbf{R}\}$. For example, $1 + 2i \in \mathbf{C}$.
- \mathbf{H} or \mathbf{H} , denoting the set of all quaternions: $\mathbf{H} = \{a + bi + cj + dk : a, b, c, d \in \mathbf{R}\}$. For example, $1 + i + 2j - k \in \mathbf{H}$.

Positive and negative sets are denoted by a superscript - or +, for example: \mathbf{Q}^+ represents the set of positive rational numbers.

Each of the above sets of numbers has an infinite number of elements, and each can be considered to be a proper subset of the sets listed below it. The primes are used less frequently than the others outside of number theory and related fields.

Basic operations

There are several fundamental operations for constructing new sets from given sets.

Unions

Main article: Union (set theory)

Two sets can be "added" together. The union of A and B , denoted by $A \cup B$, is the set of all things which are members of either A or B .

Examples:

- $\{1, 2\} \cup \{\text{red, white}\} = \{1, 2, \text{red, white}\}$.
- $\{1, 2, \text{green}\} \cup \{\text{red, white, green}\} = \{1, 2, \text{red, white, green}\}$.
- $\{1, 2\} \cup \{1, 2\} = \{1, 2\}$.

Some basic properties of unions:

- $A \cup B = B \cup A$.
- $A \cup (B \cap C) = (A \cup B) \cap C$.
- $A \subseteq (A \cup B)$.
- $A \subseteq B$ if and only if $A \cup B = B$.
- $A \cup A = A$.
- $A \cup \emptyset = A$.


Intersections

Main article: Intersection (set theory)

A new set can also be constructed by determining which members two sets have "in common". The intersection of A and B , denoted by $A \cap B$, is the set of all things which are members of both A and B . If $A \cap B = \emptyset$, then A and B are said to be *disjoint*.

Examples:

- $\{1, 2\} \cap \{\text{red, white}\} = \emptyset$.
- $\{1, 2, \text{green}\} \cap \{\text{red, white, green}\} = \{\text{green}\}$.



One of the main applications of naive set theory is constructing relations. A relation from a domain A to a codomain B is a subset of the Cartesian product $A \times B$. Given this concept, we are quick to see that the set F of all ordered pairs (x, x^2) , where x is real, is quite familiar. It has a domain set \mathbf{R} and a codomain set that is also \mathbf{R} , because the set of all squares is subset of the set of all reals. If placed in functional notation, this relation becomes $f(x) = x^2$. The reason these two are equivalent is for any given value, y that the function is defined for, its corresponding ordered pair, (y, y^2) is a member of the set F .

Axiomatic set theory

Main article: Axiomatic set theory

Although initially naive set theory, which defines a set merely as any *well-defined* collection, was well accepted, it soon ran into several obstacles. It was found that this definition spawned several paradoxes, most notably:

- Russell's paradox—It shows that the "set of all sets which do not contain themselves," i.e. the "set" $\{x : x \text{ is a set and } x \notin x\}$ does not exist.
- Cantor's paradox—It shows that "the set of all sets" cannot exist.

The reason is that the phrase *well-defined* is not very well defined. It was important to free set theory of these paradoxes because nearly all of mathematics was being redefined in terms of set theory. In an attempt to avoid these paradoxes, set theory was axiomatized based on first-order logic, and thus **axiomatic set theory** was born.

For most purposes however, naive set theory is still useful.

Principle of inclusion and exclusion

Main article: Inclusion-exclusion principle

This principle gives us the cardinality of the union of sets. $|A1 \cup A2 \cup A3 \cup A4 \cup \dots \cup An| = (|A1| + |A2| + |A3| + \dots + |An|) - (|A1 \cap A2| + |A1 \cap A3| + \dots + |An-1 \cap An|) + \dots + (-1)^{n-1} (|A1 \cap A2 \cap A3 \dots \cap An|)$

See also

- Alternative set theory
- Axiomatic set theory
- Category of sets
- Class (set theory)
- Dense set
- Family of sets
- Fuzzy set
- Internal set
- Mathematical structure
- Multiset
- Naive set theory
- Rough set
- Russell's paradox
- Scientific classification
- Set notation
- Sequence (mathematics)
- Taxonomy
- Tuple
- Boolean algebra (logic)
- Principia Mathematica

Notes

- ↑ "Eine Menge, in die Zusammenfassung bestimmter, wohlunterschiedener Objekte unserer Anschauung und unseres Denkens – welche Elemente der Menge genannt werden – zu einem Ganzen." [1] (http://www.brinkmann-da.de/math/les/les01_03.htm)
- ↑ Stoll, Robert. *Sets, Logic and Axiomatic Theories*. W. H. Freeman and Company, pp. 5.

References

- Dauben, Joseph W., *Georg Cantor: His Mathematics and Philosophy of the Infinite*, Boston: Harvard University Press (1979) ISBN 978-0-691-02447-9.
- Halmos, Paul R., *Naive Set Theory*, Princeton, N.J.: Van Nostrand (1960) ISBN 0-387-90092-6.
- Stoll, Robert R., *Set Theory and Logic*, Mineola, N.Y.: Dover Publications (1979) ISBN 0-486-63829-4.
- Velleman, Daniel, *How To Prove It: A Structured Approach*, Cambridge: University Press (2006) ISBN 978-0-521-67599-4

External links

- C2 Wiki – Examples of set operations using English operators. (<http://www.c2.com/cgi/wik/SetTheory>)

Retrieved from "http://en.wikipedia.org/w/index.php?title=Set_(mathematics)&oldid=512645842"

- $\{1, 2\} \cap \{1, 2\} = \{1, 2\}$.

Some basic properties of intersections:

- $A \cap B = B \cap A$.
- $A \cap (B \cap C) = (A \cap B) \cap C$.
- $A \cap B \subseteq A$.
- $A \cap B \subseteq B$.
- $A \cap A = A$.
- $A \cap \emptyset = \emptyset$.
- $A \subseteq B$ if and only if $A \cap B = A$.

Complements

Main article: Complement (set theory)

Two sets can also be "subtracted". The *relative complement* of B in A (also called the *set-theoretic difference* of A and B), denoted by $A \setminus B$ (or $A - B$), is the set of all elements which are members of A but not members of B . Note that it is valid to "subtract" members of a set that are not in the set, such as removing the element *green* from the set $\{1, 2, 3\}$; doing so has no effect.

In certain settings all sets under discussion are considered to be subsets of a given universal set U . In such cases, $U \setminus A$ is called the *absolute complement* or simply *complement* of A , and is denoted by A' .

Examples:

- $\{1, 2\} \setminus \{\text{red, white}\} = \{1, 2\}$.
- $\{1, 2, \text{green}\} \setminus \{\text{red, white, green}\} = \{1, 2\}$.
- $\{1, 2\} \setminus \{1, 2\} = \emptyset$.
- $\{1, 2, 3, 4\} \setminus \{1, 3\} = \{2, 4\}$.
- If U is the set of integers, E is the set of even integers, and O is the set of odd integers, then $U \setminus E = E' = O$.

Some basic properties of complements:

- $A \setminus B \neq B \setminus A$ for $A \neq B$.
- $A \cup A' = U$.
- $A \cap A' = \emptyset$.
- $(A')' = A$.
- $A \setminus A = \emptyset$.
- $U' = \emptyset$ and $\emptyset' = U$.
- $A \setminus B = A \cap B'$.

An extension of the complement is the symmetric difference, defined for sets A, B as

$$A \Delta B = (A \setminus B) \cup (B \setminus A).$$

For example, the symmetric difference of $\{7,8,9,10\}$ and $\{9,10,11,12\}$ is the set $\{7,8,11,12\}$.

Cartesian product

Main article: Cartesian product

A new set can be constructed by associating every element of one set with every element of another set. The *Cartesian product* of two sets A and B , denoted by $A \times B$ is the set of all ordered pairs (a, b) such that a is a member of A and b is a member of B .

Examples:

- $\{1, 2\} \times \{\text{red, white}\} = \{(1, \text{red}), (1, \text{white}), (2, \text{red}), (2, \text{white})\}$.
- $\{1, 2, \text{green}\} \times \{\text{red, white, green}\} = \{(1, \text{red}), (1, \text{white}), (1, \text{green}), (2, \text{red}), (2, \text{white}), (2, \text{green}), (\text{green, red}), (\text{green, white}), (\text{green, green})\}$.
- $\{1, 2\} \times \{1, 2\} = \{(1, 1), (1, 2), (2, 1), (2, 2)\}$.

Some basic properties of cartesian products:

- $A \times \emptyset = \emptyset$.
- $A \times (B \cap C) = (A \times B) \cap (A \times C)$.
- $(A \cup B) \times C = (A \times C) \cup (B \times C)$.

Let A and B be finite sets. Then

- $|A \times B| = |B \times A| = |A| \times |B|$.

Applications

Set theory is seen as the foundation from which virtually all of mathematics can be derived. For example, structures in abstract algebra, such as groups, fields and rings, are sets closed under one or more operations.

Categories: Set theory | Concepts in logic | Mathematical concepts

- This page was last modified on 15 September 2012 at 12:20.
- Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. See Terms of use for details. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.

QUIZ

1.-If $A = \{1, 5, 10, 20, 30\}$ then

- A) $10 \notin A$
- B) $10 \in A$ and $15 \in A$
- C) $10 \in A$
- D) None

QUIZ

2.-If $A = \{\text{odd numbers}\}$ and $B = \{\text{even numbers}\}$ thus $A \cup B$ is equal to:

- A) A
- B) B
- C) Set of natural numbers
- D) None

QUIZ

3.-Given $A = \{2, 4, 6, \dots\}$, $B = \{5, 10, 15, 20, \dots\}$ and $C = \{10, 20, 30, \dots\}$, then $(A \cap B) \cap C$ is equal to:

- A) A
- B) B
- C) C

- D) $A \cap C$
- E) None

- C) \emptyset
- D) None

QUIZ

4.-What is set $\{10, 20, 30\}$ of set $\{0, 10, 20, 30\}$?

- A) Equal
- B) Equivalent
- C) Subset
- D) Complement set

QUIZ

5.-Let A and B are two sets, then $A - B$ is

- A) Sub-set or equal to A
- B) Sub-set of B

QUIZ

Answers

- C)
 - D)
 - C)
 - C)
 - A)
- Do not copy :)
